

OT Mastery

Dysphagia Across the Lifespan

1. What are the two types of dysphagia?

- A. Oropharyngeal dysphagia and pharyngeal dysphagia
 - B. Sphincter dysphagia and neurological dysphagia
 - C. Esophageal dysphagia and oropharyngeal dysphagia
 - D. There is only one type of dysphagia
-

2. Esophageal dysphagia is NOT likely to cause what symptom?

- A. Unintentional weight loss
 - B. Drooling
 - C. A hoarse or weak voice
 - D. A gurgling or wet-sounding voice
-

3. What are some of the most common risk factors for dysphagia in children?

- A. Cleft palate, premature birth, dental problems, GERD
 - B. ADHD, ODD, cerebral palsy, Down syndrome
 - C. Cerebral palsy, motor delays, speech delays, selective eating habits
 - D. Autism Spectrum Disorder, families with strict mealtime routines, allergies, dietary restrictions
-

4. What type of health condition places someone at a greater risk of experiencing oropharyngeal dysphagia?

- A. Pulmonary diagnoses
 - B. Neurological disorders
 - C. Musculoskeletal conditions
 - D. Gastrointestinal diseases
-

5. What statement regarding dysphagia is true?

- A. Dysphagia is only known to impact adults and older adults.
 - B. While dysphagia can affect those of any age, the elderly are more likely to experience this concern since age is a risk factor for dysphagia.
 - C. Dysphagia can only be treated in skilled nursing facilities or outpatient facilities.
 - D. OTs are not qualified to fully treat someone with dysphagia.
-

6. What feeding-related concerns fall under an OT's scope of practice?

- A. Ventilator management, allergy testing, ruling out underlying medical conditions
 - B. Weight management, caregiver training, exercise recommendations
 - C. Cognition, diet modifications, trialing pureed foods
 - D. Upper extremity use, seating and positioning during meals and snacks, sensory concerns that impact feeding
-

7. What feeding skills might a therapist want to assess a 6-month-old child for?

- A. Sucking skills
 - B. Ability to sit up independently while eating
 - C. Interest in food and self-feeding
 - D. Utensil use
-

8. What might indicate a problem with a baby's rooting reflex?

- A. If the they don't drool at all
 - B. If their tongue sits too far back in their mouth
 - C. If they demonstrate hyposensitivity to input in and around the mouth
 - D. If it persists beyond 6 months old
-

9. What is NOT a potential cause of jaw thrust?

- A. Neck hyperextension
 - B. Hypotonia
 - C. Hypertonia
 - D. Poor seated posture
-

10. What feeding-related concern is NOT likely to result from poor posture?

- A. Jaw thrust
 - B. Tongue retraction
 - C. Exaggerated jaw movements
 - D. Jaw clenching
-

11. Stress is associated with what two clinical presentations related to feeding?

- A. Tonic bite, stability bite
 - B. Clamping the mouth shut, jaw clenching
 - C. Tongue thrust, pocketing
 - D. Jaw retraction, pocketing
-

12. What assessment tool is most appropriate for a 5-year-old child who has feeding concerns?

- A. NeoEAT
 - B. M.D. Anderson Dysphagia Index
 - C. PediEAT
 - D. Functional Oral Intake Scale
-

13. What assessment tool is most appropriate for a 1-year-old child with potential reflux?

- A. ChOMPS
 - B. NeoEAT
 - C. Neonatal Eating Outcome
 - D. GIGER Scale
-

14. What activities may be used to strengthen the cheeks?

- A. Using a straw to blow bubbles into a cup of liquid; blowing on a pinwheel; filling your mouth with air, holding, then slowly blowing out
 - B. Puckering your lips; moving the tongue side to side; making humming noises
 - C. Eating cold foods; making sounds like "AHHH" and alternating between "MABA" and "NADA"; using suction from a straw to transfer cotton balls from one place to another
 - D. Playing tug-of-war with a chewy candy; chewing on ice cubes; making silly faces in the mirror
-

15. What exercise helps raise the larynx to protect it when swallowing?

- A. Effortful swallow
 - B. Mendelsohn maneuver
 - C. Shaker exercise
 - D. Super-supraglottic swallow
-

16. What body positioning might help a 54-year-old patient who has unilateral weakness and dysphagia after a stroke?

- A. Tilting the head to the weaker side
 - B. Tilting the head to the weaker side while also doing a chin tuck
 - C. Tilting the head to the stronger side
 - D. Reclining at a 30° angle while seated
-

17. An OT might recommend an older adult avoid what foods to help manage dysphagia?

- A. Mashed potatoes
- B. Buttered biscuits
- C. Pasta in sauce

D. Crackers

18. Adaptive equipment is one of the interventions OTs can use with patients who have dysphagia. If a therapist recommends coated eating utensils, what concern might they be trying to address?

- A. Lip sensitivity or limited awareness in the oral cavity
 - B. Poor lip closure or oral motor range of motion
 - C. Need for lip and tongue stimulation or poor tongue movement
 - D. Jaw instability or inability to assume a chin tuck position
-

19. What feeding concern does a maroon spoon NOT assist with?

- A. Oral motor hypersensitivity
 - B. Tongue thrust
 - C. Jaw instability
 - D. Limited lip closure
-

20. What does AOTA NOT list as OT feeding interventions in their position statement?

- A. Environmental modifications to assist with safety and attention during mealtimes
 - B. Visual-perceptual skills to assist with accuracy of self-feeding
 - C. Non-nutritive oral stimulation, exercises, and techniques
 - D. Creating customized diets to decrease risk of aspiration in those with dysphagia
-

Copyright © 2025 OT Mastery

Visit us at <https://www.otmastery.com>